

**NOTE:** These slides contain *both* Section 1.4 and 1.5.

---

## 1.4 The Matrix Equation $A\mathbf{x} = \mathbf{b}$

---

McDonald Fall 2018, MATH 2210Q 1.4 &1.5 Slides

**1.4 Homework:** Read section and do the reading quiz. Start with practice problems, then do

- **Hand in:** 1, 3, 13, 17, 19, 22, 23, 25
- Extra Practice: 4, 7, 9, 11, 31

The definition below lets us rephrase some of the concepts from Section 1.3 by viewing linear combinations of vectors as the product of a matrix and a vector

**Definition 1.4.1.** If  $A$  is an  $m \times n$  matrix, with columns  $\mathbf{a}_1, \dots, \mathbf{a}_n$ , and  $\mathbf{x}$  is in  $\mathbb{R}^n$ , then the **product of  $A$  and  $\mathbf{x}$** , denoted by  $A\mathbf{x}$ , is the linear combination of the columns of  $A$  using the corresponding entries in  $\mathbf{x}$  as weights:

$$A\mathbf{x} = \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n.$$

**Remark 1.4.2.**  $A\mathbf{x}$  is only defined the number of columns of  $A$  equals the number of entries in  $\mathbf{x}$ .

**Example 1.4.3.** Find the following products:

(a)  $\begin{bmatrix} 1 & 2 & -1 \\ 0 & -5 & 3 \end{bmatrix} \begin{bmatrix} 4 \\ 3 \\ 7 \end{bmatrix}$

(b)  $\begin{bmatrix} 1 & 2 & 1 \\ 3 & 9 & 6 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix}$

**Example 1.4.4.** Compute  $A\mathbf{x}$ , where  $A = \begin{bmatrix} 2 & 3 & 4 \\ -1 & 5 & -3 \\ 6 & -2 & 8 \end{bmatrix}$  and  $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$

**Procedure 1.4.5** (Row Vector Rule). If  $A\mathbf{x}$  is defined, then the  $i$ th entry in  $A\mathbf{x}$  is the sum of the products of corresponding entries from row  $i$  of  $A$  and from the vector  $\mathbf{x}$ .

**Example 1.4.6.** Compute

(a)  $\begin{bmatrix} 2 & -3 \\ 8 & 0 \\ -5 & 2 \end{bmatrix} \begin{bmatrix} 4 \\ 7 \end{bmatrix}$

(b)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix}$

**Example 1.4.7.** Write the system below as  $A\mathbf{x} = \mathbf{b}$  for some  $A$  and  $\mathbf{b}$ .

$$x_1 + 2x_2 - x_3 = 4$$

$$-5x_2 + 3x_3 = 1$$

**Definition 1.4.8.** The equation  $A\mathbf{x} = \mathbf{b}$  is called a **matrix equation**.

**Theorem 1.4.9.** If  $A$  is an  $m \times n$  matrix, with columns  $\mathbf{a}_1, \dots, \mathbf{a}_n$ , and  $\mathbf{b}$  is in  $\mathbb{R}^m$ , then

$$A\mathbf{x} = \mathbf{b} \text{ (with } \mathbf{x} \text{ in } \mathbb{R}^n \text{)}$$

has the same solutions as the vector equation

$$x_1\mathbf{a}_1 + x_2\mathbf{a}_2 + \cdots + x_n\mathbf{a}_n = \mathbf{b}$$

which has the same solutions as the system of linear equations with augmented matrix

$$\left[ \begin{array}{cccc|c} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n & \mathbf{b} \end{array} \right].$$

**Corollary 1.4.10.** The equation  $A\mathbf{x} = \mathbf{b}$  has a solution if and only if  $\mathbf{b}$  is a linear combination of the columns of  $A$ .

**Example 1.4.11.** Let  $A = \begin{bmatrix} 1 & 3 & 4 \\ -4 & 2 & -6 \\ -3 & -2 & -7 \end{bmatrix}$  and  $\mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$  is  $A\mathbf{x} = \mathbf{b}$  consistent for all  $b_1, b_2, b_3$ ?

**Theorem 1.4.12.** *Let  $A$  be an  $m \times n$  matrix. Then the following statements are either all true, or all false.*

- (a) *For each  $\mathbf{b}$  in  $\mathbb{R}^m$ , the equation  $A\mathbf{x} = \mathbf{b}$  has a solution.*
- (b) *Each  $\mathbf{b}$  in  $\mathbb{R}^m$  is a linear combination of the columns of  $A$ .*
- (c) *The columns of  $A$  span  $\mathbb{R}^m$ .*
- (d)  *$A$  has a pivot position in every row.*

**Example 1.4.13.** If  $A = \begin{bmatrix} 2 & 0 & -2 \\ 2 & 3 & 4 \\ 0 & 1 & 2 \end{bmatrix}$  and  $\mathbf{b} = \begin{bmatrix} 2 \\ 11 \\ 3 \end{bmatrix}$ , for what  $\mathbf{x}$  is  $A\mathbf{x} = \mathbf{b}$  consistent?

We end this section with some important properties of  $A\mathbf{x}$ , which we will use throughout the course.

**Theorem 1.4.14.** *If  $A$  is an  $m \times n$  matrix,  $\mathbf{u}$  and  $\mathbf{v}$  are vectors in  $\mathbb{R}^n$ , and  $c$  is a scalar:*

- (a)  $A(\mathbf{u} + \mathbf{v}) = A\mathbf{u} + A\mathbf{v}$ ;
- (b)  $A(c\mathbf{u}) = c(A\mathbf{u})$ .

---

## 1.5 Solutions Sets of Linear Systems

---

**1.5 Homework:** Read section and do the reading quiz. Start with practice problems, then do

- **Hand in:** 5, 11, 15, 19, 23, 30, 32
- Extra Practice: 2, 6, 18, 22, 27

In this section, we will use vector notation to give explicit and geometric descriptions of solution sets of linear systems. We begin by defining a special type of system.

**Definition 1.5.1.** A system of linear equations is said to be **homogeneous** if it can be written in the form  $A\mathbf{x} = \mathbf{0}$ , where  $A$  is an  $m \times n$  matrix, and  $\mathbf{0}$  is the zero vector in  $\mathbb{R}^m$ .

**Remark 1.5.2.** The equation  $A\mathbf{x} = \mathbf{0}$  *always* has at least one solution, namely  $\mathbf{x} = \mathbf{0}$ , called the **trivial solution**. We will be interested in finding **non-trivial solutions**, where  $\mathbf{x} \neq \mathbf{0}$ .

**Example 1.5.3.** Determine if the following homogeneous system has a nontrivial solution, and describe the solution set.

$$\begin{aligned}3x_1 + 5x_2 - 4x_3 &= 0 \\-3x_1 - 2x_2 + 4x_3 &= 0 \\6x_1 + x_2 - 8x_3 &= 0\end{aligned}$$

**Proposition 1.5.4.** *The homogeneous equation  $A\mathbf{x} = \mathbf{0}$  has a nontrivial solution if and only if the equation has at least one free variable.*

**Example 1.5.5.** Describe all solutions to the homogeneous system

$$10x_1 - 3x_2 - 2x_3 = 0.$$

**Definition 1.5.6.** The answers in 1.5.3 and 1.5.5 are **parametric vector equations**. Sometimes, to emphasize that the parameters vary over all real numbers, we write


$$\mathbf{x} = s\mathbf{u} + t\mathbf{v} \text{ for } s, t \in \mathbb{R}.$$

In both examples, we say that the solution is in **parametric vector form**.

**Example 1.5.7.** Describe all solutions of

$$\begin{aligned} 3x_1 + 5x_2 - 4x_3 &= 7 \\ -3x_1 - 2x_2 + 4x_3 &= -1 \\ 6x_1 + x_2 - 8x_3 &= -4 \end{aligned}$$


**Definition 1.5.8.** We can think of vector addition as *translation*. Given  $\mathbf{p}$  and  $\mathbf{v}$  in  $\mathbb{R}^2$  or  $\mathbb{R}^3$ , the effect of adding  $\mathbf{p}$  to  $\mathbf{v}$  is to *move*  $v$  in a direction parallel to the line through  $\mathbf{p}$  and  $\mathbf{0}$ . We say that  $\mathbf{v}$  is **translated by  $\mathbf{p}$**  to  $\mathbf{v} + \mathbf{p}$ . If each point on a line  $L$  is translated by a vector  $\mathbf{p}$ , the result is a line parallel to  $L$ .



For  $t \in \mathbb{R}$ , we call  $\mathbf{p} + t\mathbf{v}$  **the equation of the line parallel to  $\mathbf{v}$  through  $\mathbf{p}$** .

**Example 1.5.9.** Use this observation to describe the relationships between the solutions to  $A\mathbf{x} = \mathbf{0}$  and  $A\mathbf{x} = \mathbf{b}$  using the  $A$  and  $\mathbf{b}$  from Examples 1.5.3 and 1.5.7.

**Theorem 1.5.10.** Suppose the equation  $A\mathbf{x} = \mathbf{b}$  is consistent for some given  $\mathbf{b}$ , and let  $\mathbf{p}$  be a solution. Then the solution set of  $A\mathbf{x} = \mathbf{b}$  is the set of all vectors of the form  $\mathbf{w} = \mathbf{p} + \mathbf{v}_h$ , where  $\mathbf{v}_h$  is any solution of the homogeneous equation  $A\mathbf{x} = \mathbf{0}$ .



**Procedure 1.5.11.** To write a solution set in parametric vector form

1. Row reduce the augmented matrix to RREF
2. Express each basic variable in terms of any free variables
3. Write  $\mathbf{x}$  as a vector whose entries depend on the free variables (if there are any)
4. Decompose  $\mathbf{x}$  into a linear combination of vectors using free variables as parameters

**Example 1.5.12.** Describe and compare the solution sets of  $A\mathbf{x} = \mathbf{b}$  and  $A\mathbf{x} = \mathbf{0}$  if

$$A = \begin{bmatrix} 1 & 3 & -5 \\ 1 & 4 & -8 \\ -3 & -7 & 9 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} 4 \\ 7 \\ -6 \end{bmatrix}.$$